

# Spin-selective transport phenomena in helical molecular wires

R. Korytár\*, Š. Marek\*, J. van Ruitenbeek<sup>‡</sup> and F. Evers<sup>†</sup>

\* Charles University, Czech Republic, e-mail: korytar@karlov.mff.cuni.cz

<sup>†</sup> Institute of Theoretical Physics, University of Regensburg, Germany

<sup>‡</sup> Huygens-Kamerlingh Onnes Laboratory, Leiden University, Netherlands

## Background

Recently various spin-selective phenomena have been reported in transport through helical molecular wires attached to metallic leads[1], [2], [3], see Fig. 1. There is a general consensus that the spin selectivity results due to a combination of the spin-orbit coupling and the helicity. Theoretical understanding of the so called chirality-induced spin selectivity (CISS) is, however, not complete, because most theoretical reports grossly underestimate the magnitude of the effect [4].

We present two approaches to improve the quantitative description of the CISS.

### 1. Metal-molecular interfaces in the *GW* approximation

First-principles quantum transport simulations rely on an accurate description of electronic charged excitations. The latter are not well described in density functional theory; a natural framework to describe them is Hedin's *GW* approximation. Our calculations show that the *GW* method widens the HOMO-LUMO gap of metallic clusters. We rigorously quantify this observation by investigating an ensemble of disordered metallic clusters [6], see Fig. 2. Our observation has profound impact on the interpretation of standard calculations of molecular adsorbates. Namely, in such studies the infinite metallic surface is always replaced by a finite cluster (with periodic or vacuum boundary conditions). Our results imply that the density of states from the *GW* quasiparticles will be artificially depleted right at the Fermi level.

Therefore, the *GW* method can not be taken as an out-of-the-box improvement over the less

accurate density-functional theory.

### 2. Spin currents in chiral molecular junctions

Considerations based on time-reversal invariance known in the field of spintronics allow for spin-currents in two-terminal coherent devices even in absence of external magnetic fields and magnetic impurities. Such spin-currents exist even in linear response if the device hosts more than one scattering channel. We exemplify this in a model of a molecular wire of a helical topology, see Fig. 3 for an example. We provide basic principles for designing molecular junctions that generate sizable spin currents [5]. These principles can be verified by ab-initio calculations and experimental setups.

## Acknowledgment

R. Korytár acknowledges the Czech Science foundation (project no. 22-22419S).

## References

- [1] S. Mishra, A. Kumar Mondal, S. Pal, T. Kumar Das, E. Z. B. Smolinsky, G. Siligardi, and R. Naaman, *J. Phys. Chem. C* 124, 10776 (2020)
- [2] T. Liu, X. Wang, H. Wang, G. Shi, F. Gao, H. Feng, H. Deng, L. Hu, E. Lochner, P. Schlottmann, S. von Molnár, Y. Li, J. Zhao, and P. Xiong, *ACS Nano* 14, 15983 (2020)
- [3] H. Al-Bustami, S. Khaldi, O. Shoseyov, S. Yochelis, K. Killi, I. Berg, E. Gross, Y. Paltiel, and R. Yerushalmi, *Nano Lett.* 22, 5022 (2022)
- [4] F. Evers, et al., *Adv. Mater.* 34(13), 2106629 (2022)
- [5] J. van Ruitenbeek, R. Korytár, F. Evers, Chirality-controlled spin scattering through quantum interference, in preparation (2023)
- [6] Š. Marek and R. Korytár, Pitfalls of ab-initio *GW* calculations for metal-molecular interfaces, in preparation (2023)




Fig. 1. Typical setup to measure the CISS in electronic current: a non-magnetic lead (Au) attaches to a helical (DNA-like) molecule. The spin selectivity manifests as the the magnetization of the right lead (a ferromagnet) reverses.




Fig. 2. Scaling of various charged excitations of metallic clusters in  $evGW$ . Top: average level spacings and gaps in an ensemble of metallic clusters as a function of the number of atoms. Lines are guides to the eye; they reveal that the HOMO-LUMO gaps scale with a slower power law, leading to HOMO-LUMO widening. Right, the energy levels of sample clusters is shown, with HOMO aligned to 0.




Fig. 3. Charge conductance  $G$  and spin left/right spin conductances  $G_z^{(L,R)}$  of a model helical wire with 3 (6) turns are depicted by solid (dashed) lines as a function of the Fermi energy. The helix is attached to non-helical leads.